













- Johnston JS, Bennett MD, Rayburn AL, Galbraith DW, Price HJ (1999), 'Reference standards for determination of DNA content of plant nuclei,' *Am J Bot*, 86:609–613.
- Signor CL, Gallardo K, Prospero JM, Salon C, Quillien L, Thompson R and Duc G. (2005), 'Genetic diversity for seed protein composition in *Medicago truncatula*,' *Plant Genetic Resources*, 3(1):59–71.
- Sammour R, Mustafa A, Badr S and Taher W. (2007), 'Genetic variations in accessions of *Lathyrus sativus* L.,' *Acta Bot. Croat.* 66 (1): 1–13.
- Ochatt S, Sangwan R, (2008), 'In vitro shortening of generation time in *Arabidopsis thaliana*,' *Plant Cell Tissue Organ Cult*, 93:133–137.
- Agayev YM (1998), 'Advanced squash methods for investigation of plant chromosomes,' *Journal of Heredity*, 79(4):225-238.
- Laemmli UK (1970), 'Cleavage of structural proteins during the assembly of the head of bacteriophage T4,2 *Nature*, 227:680–685.
- Cooper, J. W., Wilson, M. H., Derks, M. F. L., Smit, S., Kunert, K. J., Cullis, C., & Foyer, C. H. (2017), 'Enhancing faba bean (*Vicia faba* L.) genome resources,' *Journal of Experimental Botany*, 68(8), 1941-1953. <https://doi.org/10.1093/jxb/erx117>
- Kubaláková, M., Valárik, M., Bartoš, J., Vrána, J., Číhalíková, J., Molnár-Lang, M., Doležel, J.: (2003), 'Analysis and sorting of rye (*Secale cereale* L.) chromosomes using flow cytometry,' *Genome* 46: 893-905.